

Observation of bosonic coalescence of photon pairs

Giovanni Di Giuseppe,^{1*} Mete Atatüre,^{2†} Matthew D. Shaw,^{1‡}
Alexander V. Sergienko,^{1,2} Bahaa E. A. Saleh,¹ Malvin C. Teich,^{1,2}

¹*Department of Electrical & Computer Engineering, and*

²*Department of Physics,*

*Quantum Imaging Laboratory,[§] Boston University,
8 Saint Mary's Street, Boston, MA 02215*

Aaron J. Miller, Sae Woo Nam, and John Martinis

*National Institute of Standards and Technology[¶],
Mail Code 814, 325 Broadway, Boulder, CO 80395*

(Dated: February 1, 2008)

Quantum theory predicts that two indistinguishable photons incident on a beam-splitter interferometer stick together as they exit the device (the pair emerges randomly from one port or the other). We use a special photon-number-resolving energy detector for a direct loophole-free observation of this quantum-interference phenomenon. Simultaneous measurements from two such detectors, one at each beam-splitter output port, confirm the absence of cross-coincidences.

PACS numbers: 42.50.Dv, 42.50.Xa, 42.50.St

Introduction.—The seminal experiment carried out by Hong, Ou, and Mandel some fifteen years ago [1] is one of the most important in the annals of quantum optics. This experiment demonstrated that two indistinguishable photons, incident on the two ports of a simple beam splitter, interfere in such a way as to always emerge as a pair, exiting randomly from one port or the other. The photons stuck together, so to speak. They could only observe this phenomenon indirectly, however, since traditional single-photon-counting detectors cannot register more than one photon within the dead-time period of the device. Hong, Ou, and Mandel circumvented this technical limitation by designing an experiment in which they measured the complement of what they sought to observe. They used two single-photon-counting detectors, placing one at each output port of the beam splitter, and then searched for cross-coincidences between the two detectors. Finding none, they inferred that the two photons do not exit from different ports of the beam splitter. Based on this “test of exclusion”, and energy conservation, they deduced that both photons must exit via the same port of the device. There have been many variations on this original theme but all rely on the same “test of exclusion”, whatever the source of the single photons [2, 3, 4, 5].

We have carried out a polarization version of the Hong-Ou-Mandel experiment and report a *direct loophole-free*

observation of photon coincidences at a single output port of the interferometer, thereby confirming the bosonic coalescence of pairs of photons in a single mode [3]. What makes this possible is a unique energy detector [6] that has the capability of registering the *number* of photons impinging on it. Our experimental results serve to provide a full confirmation of the quantum theory of photon interference in a beam-splitter interferometer [7, 8, 9, 10].

We have also made measurements at both output ports of the beam splitter using two such detectors. This has permitted us to demonstrate the enhancement of coincidences at a single output port concomitantly with the diminution of cross-coincidences at the two output ports. We thus simultaneously conduct two experiments: a new single-port photon coincidence measurement and the original photon cross-coincidence measurement. Our observations confirm the inferences made by Hong, Ou, and Mandel [1] and Shih and Sergienko [11].

Experiment.—The experimental arrangement, shown in Fig. 1, is similar to that used in related quantum-interference experiments [12], but photon-number resolving (PNR) transition-edge sensors [6] replace the usual photon-counting avalanche photodiodes.

To generate orthogonally polarized photon pairs, a single-line 351.1-nm continuous-wave (cw) argon-ion laser operated at 100 mW was used to pump a 0.5-mm-thick β -BaB₂O₄ (BBO) nonlinear-optical crystal, aligned for type-II collinear degenerate spontaneous parametric down-conversion (SPDC). The low power of the pump ensures that the photon pairs are well separated in time. A 2-mm aperture placed 70 cm beyond the crystal was used to select only those photon pairs that propagate collinearly with the pump. The pump is disposed of by use of a dichroic mirror and an RG-630 colored glass filter.

At the heart of the interferometer is a non-polarizing beam splitter that distributes the photon pairs into two

*Also at: Istituto Elettrotecnico Nazionale G. Ferraris, Strada delle Cacce 91, I-10153 Torino, Italy.

†Present Address: Institute of Quantum Electronics, ETH Hönggerberg HPT, CH-8093 Zürich, Switzerland.

‡Present Address: Stanford Linear Accelerator Center, 2575 Sand Hill Road, Menlo Park, California 94025, USA.

§URL: <http://www.bu.edu/qil>

¶Contribution of the U.S. Government; not subject to copyright.




FIG. 1: Schematic of the polarization-based analog of the Hong-Ou-Mandel interference experiment using type-II collinear degenerate spontaneous parametric down-conversion. Photon-number resolving (PNR) detectors permit the direct observation of photon coincidences at a single output port of the beam-splitter interferometer as well as cross-coincidences at the two output ports.

spatial modes, denoted A and B in Fig. 1. In each arm, a Glan-Thompson polarization analyzer oriented at 45° renders the two photons indistinguishable in polarization. Narrowband interference filters were not used in the experiment.

A lens in each arm (not shown) couples the light into two $9\text{-}\mu\text{m}$ mode-field-diameter optical fibers, each of which is connected to a separate PNR detector operated in a cryostat. Detection events from the two PNR detectors were recorded and analyzed using a computer-controlled system that registers time-stamped single- and double-photon events. Software is used to extract the coincidences at a single detector and the cross-coincidences from the two detectors.

The detector elements of the transition-edge sensor are photolithographically patterned 40-nm-thick tungsten thin films deposited on a silicon substrate [6]. The substrate is cooled to approximately 60 mK, about half the superconducting-to-normal transition temperature of 100 mK (the transition width is about 1 mK). A bias voltage across the thin film maintains the temperature in the transition region via Joule heating. An incident photon absorbed by the tungsten film is converted to a photoelectron, which raises the electron temperature of the film, thereby increasing its resistance. The time integral of the associated decrease in current, multiplied by the bias voltage, provides the total photoelectric energy absorbed by the thin film within its $15\text{-}\mu\text{sec}$ thermal relaxation time. In conducting an optical experiment using light of a specified wavelength, the number of photons incident within the thermal relaxation time is determined

by establishing the total energy transferred to the detector within this time. Of course, energy detectors of this kind cannot distinguish between the absorption of two photons, each of energy E , and the absorption of a single photon of energy $2E$; great care was therefore used to prevent UV pump photons from leaking through to the detectors.

Over the range of wavelengths of interest in our experiments, the quantum efficiency η of these PNR detectors is approximately 20% [13], as determined via an absolute measurement technique [14, 15, 16]. The finite quantum efficiency obviates the possibility of determining the incident photon number with certainty. Nevertheless, by virtue of the invariance of the Poisson distribution to random deletion [17], it is possible to infer the photon-number resolving capability of these detectors by measuring the number distribution that results from excitation with weak laser light [13]. The typical full-width half-maximum energy resolution of these detectors is currently about 0.25 eV at 1.77 eV, corresponding to 100 nm at a central wavelength of 700 nm. The signal is read out of the detector using a system that incorporates an array of dc superconducting quantum interference devices (SQUIDS), which operate as current-sensitive amplifiers. To limit spurious “pileup” counts, experiments are carried out using a reduced singles counting rate ~ 1000 counts/sec, an order of magnitude lower than the achievable counting rate.

A Babinet-type compensator consisting of two parallel quartz prisms and a fixed quartz plate, z -cut to eliminate transverse birefringence, is used to modify the relative time delay τ between the two photons. The condition of indistinguishability is imposed by overlapping the two single-photon wave packets in space and time at the beam splitter; this is not required, in principle, as long as distinguishing information is subsequently erased [18].

An experiment is conducted by modifying the degree of distinguishability (achieved by varying the delay time τ over a range of hundreds of fsec) and tracing out three curves: the coincidence probability at detector A, denoted $P(2, 0)$; the coincidence probability at detector B, denoted $P(0, 2)$; and the cross-coincidence probability, denoted $P(1, 1)$.

The results of a typical experiment are illustrated in Fig. 2. Never-before seen are the data for the peaks in the coincidence probabilities, $P(2, 0)$ and $P(0, 2)$, at detectors A and B, respectively. These peaks are a manifestation of excess photon pairs at a single output port of the beam splitter. The coincidence probability is enhanced because the photons are identical bosons when the wave packets overlap. Were the same experiment to be conducted with fermions, the coincidence probabilities would be suppressed rather than enhanced [19]. The data points for the complementary cross-coincidence probability $P(1, 1)$ (diamonds) exhibit the familiar dip associated with the quantum interference of indistinguishable photons [12].

Delay times τ that are substantially larger, or smaller,




FIG. 2: Experimental coincidence probabilities $P(2,0)$ and $P(0,2)$, and cross-coincidence probability $P(1,1)$, as a function of the relative delay time between the photons τ imparted by the compensator (symbols). The theoretical curves (dashed) are computed from the ideal single-mode theory provided in Eqs. (3) and (4), assuming a crystal of length 0.5 mm. The data follow the trends of the theory well, but with reduced visibility resulting from imperfect alignment and asymmetric polarization losses in the system.

than those at the dip/peak correspond to distinguishable photons. In that domain the coincidence probabilities are characterized by classical particle-like statistics, namely the binomial counting distribution [9]. The cross-coincidence probability is then expected to be twice the coincidence probability, which is, in fact, exactly what is observed on the shoulders of the interference pattern (see Fig. 2).

Theory.—For collinear SPDC confined to a single spatial mode, the quantum state at the output of the nonlinear crystal driven by a monochromatic plane-wave pump at frequency ω_p , is [12]

$$|\psi\rangle = \int d\omega \tilde{\Phi}(\omega) \hat{a}_o^\dagger\left(\frac{\omega_p}{2} + \omega\right) \hat{a}_e^\dagger\left(\frac{\omega_p}{2} - \omega\right) |0\rangle \quad (1)$$

where the operators $\hat{a}_o^\dagger(\omega)$ and $\hat{a}_e^\dagger(\omega)$ create photons of ordinary and extraordinary polarization, respectively, in frequency mode ω , and the limits on the integral stretch from $-\infty$ to ∞ . The form assumed by the state function $\tilde{\Phi}(\omega)$, which is normalized according to $\int d\omega |\tilde{\Phi}(\omega)|^2 = 1$ depends on the physical structure of the down-conversion source. The state function for a single bulk crystal of length L is, for example, given by [12]

$$\tilde{\Phi}(\omega) \propto L \operatorname{sinc}\left[\frac{L}{2} \Delta(\omega)\right] e^{iL\Delta(\omega)/2}, \quad (2)$$

where the wave-vector mismatch function $\Delta(\omega) = k_p(\omega_p) - k_o(\omega_p/2 + \omega) - k_e(\omega_p/2 - \omega)$ depends on the dispersive properties of the birefringent medium ($k_{\{p,o,e\}}$

represent the wave numbers of the pump, ordinary wave, and extraordinary wave in the crystal, respectively).

Assuming an ideal optical system and perfect components, the coincidence probabilities, as a function of τ , are then given by

$$P(2,0) = P(0,2) = \frac{1}{32} \int dt |\Phi(t - \tau) + \Phi(-t - \tau)|^2, \quad (3)$$

where $\Phi(t)$ is the inverse Fourier transform of $\tilde{\Phi}(\omega)$ and the integration can be extended from $-\infty$ to ∞ since the integrand is narrow in comparison with the detection window. Similarly, the cross-coincidence probability becomes [20]

$$P(1,1) = \frac{1}{16} \int dt |\Phi(t - \tau) - \Phi(-t - \tau)|^2. \quad (4)$$

The fractional pre-factors in Eqs. (3) and (4), $1/32$ and $1/16$ respectively, can be traced to the fact that the polarization analyzers in each arm of the interferometer transmit only half the photons.

For perfect photon-wavepacket overlap, with the value of τ chosen such that $\Phi(t - \tau) = \Phi(-t - \tau)$, Eq. (3) becomes $P(2,0) = P(0,2) = 2^2/32 = 1/8$ while Eq. (4) becomes $P(1,1) = 0$, and the result reduces to the ideal boson counting distribution.

In the opposite limit, when the relative delay time τ is sufficiently large, the integrals in Eqs. (3) and (4) are both equal to 2, whereupon the classical binomial counting distribution emerges: $P(2,0) = P(0,2) = 1/16$ and $P(1,1) = 1/8$.

More generally, Eqs. (3) and (4) can be combined to provide a complementary formula that relates the coincidence and cross-coincidence probabilities for arbitrary values of τ :

$$P(1,1) + P(2,0) + P(0,2) = \frac{1}{4}. \quad (5)$$

In particular, when the state function in Eq. (2) is used in Eqs. (3) and (4), assuming a cw pump field and a linear approximation for $\Delta(\omega)$, the cross-coincidence probability $P(1,1)$, as a function of τ , takes the familiar form of a triangular dip [11, 12] while the coincidence probabilities $P(2,0)$ and $P(0,2)$ behave as triangular peaks.

The theoretical results for this ideal single-mode theory are shown as the dashed curves in Fig. 2. The data follow the trends of the theory well though the visibilities of the quantum interference patterns are reduced below their ideal values as a result of imperfect alignment and asymmetric polarization losses in the optical components.

Conclusion.—We have used special photon-number-resolving energy detectors to directly demonstrate that two indistinguishable photons incident on a polarization analog of the Hong-Ou-Mandel interferometer stick together as they exit the beam-splitter ports. The absence of cross-coincidences has also been concomitantly demonstrated. Our observations provide a full confirmation of

the quantum-optical theory of interference in a beam-splitter interferometer. It is expected that similar results would be observed for a non-collinear configuration. As a final note we point out that PNR detectors, such as those used here, are expected to find use in other quantum-optics and quantum-information-processing experiments. Unlike their APD counterparts, they could play a role in carrying out conclusive tests of local realism using a beam-splitter interferometer experiment [21]. They have

already been found to be useful in a number of other important applications [22].

Acknowledgments. — We are grateful to A. Abouraddy for valuable comments. This work was supported by the National Science Foundation; the Center for Subsurface Sensing and Imaging Systems (CenSSIS), an NSF Engineering Research Center; the Defense Advanced Research Projects Agency (DARPA); and the David & Lucile Packard Foundation.

-
- [1] C. K. Hong, Z. Y. Ou, and L. Mandel, *Phys. Rev. Lett.* **59**, 2044 (1987).
 - [2] J. G. Rarity and P. R. Tapster, *J. Opt. Soc. Am. B* **6**, 1221 (1989).
 - [3] P. Grangier, *Nature* **419**, 577 (2002).
 - [4] C. Santori, D. Fattal, J. Vučković, G. S. Solomon, and Y. Yamamoto, *Nature* **419**, 594 (2002).
 - [5] H. Kim, J. Ko, and T. Kim, *J. Opt. Soc. Am. B* **20**, 760 (2003).
 - [6] B. Cabrera, R. M. Clarke, P. Colling, A. J. Miller, S. Nam, and R. W. Romani, *Appl. Phys. Lett.* **73**, 735 (1998).
 - [7] B. Yurke, S. L. McCall, and J. R. Klauder, *Phys. Rev. A* **33**, 4033 (1986).
 - [8] H. Fearn and R. Loudon, *J. Opt. Soc. Am. B* **6**, 917 (1989).
 - [9] R. A. Campos, B. E. A. Saleh, and M. C. Teich, *Phys. Rev. A* **40**, 1371 (1989).
 - [10] R. A. Campos, B. E. A. Saleh, and M. C. Teich, *Phys. Rev. A* **42**, 4127 (1990).
 - [11] Y. H. Shih and A. V. Sergienko, *Phys. Lett. A* **186**, 29 (1994).
 - [12] M. Atatüre, G. Di Giuseppe, M. D. Shaw, A. V. Sergienko, B. E. A. Saleh, and M. C. Teich, *Phys. Rev. A* **66**, 023822 (2002).
 - [13] A. J. Miller, S. W. Nam, J. M. Martinis, and A. V. Sergienko, Submitted to *Appl. Phys. Lett.* (2003).
 - [14] D. N. Klyshko, *Kvantovaya Elektron.* **7**, 1932 (1980) [Translation: *Sov. J. Quantum Electron.* **10**, 1112 (1980)].
 - [15] J. G. Rarity, K. D. Ridley, and P. R. Tapster, *Appl. Opt.* **26**, 4616 (1987).
 - [16] A. Czitrovsky, A. V. Sergienko, P. Jani, and A. Nagy, *Metrologia* **37**, 617 (2000).
 - [17] M. C. Teich and B. E. A. Saleh, *Optics Lett.* **7**, 365 (1982).
 - [18] T. B. Pittman, D. V. Strekalov, A. Migdall, M. H. Rubin, A. V. Sergienko, and Y. H. Shih, *Phys. Rev. Lett.* **77**, 1917 (1996).
 - [19] R. Loudon, *Phys. Rev. A* **58**, 4904 (1998).
 - [20] M. H. Rubin, D. N. Klyshko, Y. H. Shih, and A. V. Sergienko, *Phys. Rev. A* **50**, 5122 (1994).
 - [21] S. Popescu, L. Hardy, and M. Zukowski, *Phys. Rev. A* **56**, R4353 (1997).
 - [22] R. W. Romani, A. J. Miller, B. Cabrera, S. W. Nam, and J. M. Martinis, *Astrophys. J.* **563**, 221 (2001).